## WHAT IS CLAIMED IS:

1. A method of carrying out a reaction comprising the steps of:

mixing at least one organic reaction component with a fluorous reaction component having the formula:

 $X^{1}Sn(R)_{n}[Rs(Rf)]_{3-n}, X^{1}X^{2}Sn[Rs(Rf)]_{2} \text{ or } O=Sn[Rs(Rf)]_{2}$ 

wherein n is 1 or 2, R is a  $C_1$ - $C_6$  alkyl group,  $X^1$  and  $X^2$  are independently, the same or different, H, F, Cl, Br, I,  $N_3$ ,  $OR^1$ ,  $OOR^1$ ,  $SR^1$ ,  $SeR^1$ , CN, NC,  $NR^1R^2$ , an aryl group, a heteroaryl group, an alkyl group of 1 to 20 carbons, an alkenyl group, an alkynyl group,  $-C(O)R^3$ ,  $M((Rs')(Rf'))_3$ ,  $OM((Rs')(Rf'))_3$  or  $OOM((Rs')Rf'))_3$ , wherein M is Si, Ge, or Sn, and wherein  $R^1$  and  $R^2$  are each independently the same or different H, an alkyl group,  $-SO_2R^3$  or  $-C(O)R^3$ , wherein  $R^3$  is an alkyl group or an aryl group, and wherein Rs and Rs' are each independently the same or different a spacer group, and wherein Rf and Rf' are each independently the same or different a fluorous group;

carrying out a reaction to produce an organic product; and

after producing the organic product, separating any excess of the fluorous reaction component and any fluorous byproduct of the fluorous reaction component using a fluorous separation technique.

- 2. The method of Claim 1 wherein  $X^1$  and  $X^2$  are independently the same or different an allyl group, Br, Cl, F, I, or H, Rs is  $-CH_2CH_2-$ , and Rf is a perfluoroalkyl group.
- 3. The method of Claim 1 wherein Rf is a fluorohydrocarbon group, a perfluorocarbon group, a fluorinated ether group or a fluorinated amine group.
- 4. The method of Claim 1 wherein Rf is a linear perfluoroalkyl group of 3 to 20 carbons, a branched perfluoroalkyl group of 3 to 20 carbons, and a hydrofluoroalkyl group of 3 to 20 carbons, the hydrofluoroalkyl group comprising up to one hydrogen atom for each two fluorine atoms.
- 5. The method of Claim 1 wherein Rf is a linear perfluoroalkyl group of 6 to 12 carbons, a branched perfluoroalkyl group of 6 to 12 carbons, or a hydrofluoroalkyl group of 6 to 12 carbons, the hydrofluoroalkyl group comprising up to one hydrogen atom for each two fluorine atoms.
- 6. The method of Claim 1 wherein R<sup>3</sup> is a perfluoroalkyl group.
- 7. The method of Claim 1 wherein Rs is an alkylene group or a phenylene group.
- 8. The method of Claim 1 wherein Rs is an alkylene group.

9. A method for carrying out a chemical reaction, comprising the steps of:

combining at least one fluorous reaction component having the formula:

 $X^{1}Sn(R)_{n}[Rs(Rf)]_{3-n}, X^{1}X^{2}Sn[Rs(Rf)]_{2} \text{ or } O=Sn[Rs(Rf)]_{2}$ 

wherein n is 1 or 2, R is a  $C_1$ - $C_6$  alkyl group,  $X^1$  and  $X^2$  are independently, the same or different, H, F, Cl, Br, I,  $N_3$ ,  $OR^1$ ,  $OOR^1$   $SR^1$ ,  $SeR^1$ , CN, NC,  $NR^1R^2$ , an aryl group, a heteroaryl group, an alkyl group of 1 to 20 carbons, an alkenyl group, an alkynyl group,  $-C(O)R^3$ ,  $M((Rs')(Rf'))_3$ ,  $OM((Rs')(Rf'))_3$  or  $OOM((Rs')Rf'))_3$ , wherein M is Si, Ge, or Sn, and wherein  $R^1$  and  $R^2$  are each independently the same or different H, an alkyl group,  $-SO_2R^3$  or  $-C(O)R^3$ , wherein  $R^3$  is an alkyl group or an aryl group, and wherein Rs and Rs' are each independently the same or different a spacer group, and wherein Rf and Rf' are each independently the same or different a fluorous group, and at least one organic reaction component convertible in the presence of the fluorous reaction component to a product in an organic solvent;

contacting the fluorous reaction component and the organic reaction component in the organic solvent under conditions suitable to produce the product; and

after production of the product, separating any excess of the fluorous reaction component and any fluorous byproduct of the fluorous reaction component using a fluorous separation technique. put 63

 $\sqrt{10}$ . A chemical compound having the formula:

 $X^{1}Sn(R)_{n}[Rs(Rf)]_{3-n}$ 

wherein n is 1 or  $\chi$ , R is a  $C_1$ - $C_6$  alkyl group,  $\chi^1$  is H, F, Cl, Br, I,  $N_3$ ,  $OR^1$ ,  $OR^1$   $SR^1$ ,  $SeR^1$ , CN, NC,  $NR^1R^2$ , an aryl group, a heteroaryl group, an alkyl group of 1 to 20 carbons, an alkenyl group, an alkynyl group,  $-C(0)R^3$ ,  $M((Rs')(Rf'))_3$ ,  $OM((Rs')(Rf'))_3$  or  $OOM((Rs')Rf'))_3$ , wherein M is Si, Ge, or Sn, and  $\backslash$  wherein  $R^1$  and  $R^2$  are each independently the same or different H, an alkyl group,  $-SO_2R^3$  or  $-C(0)R^3$ , wherein  $R^3$  is an alkyl group or an aryl group, and wherein Rs and Rs' are each independently the same or different an alkylene group of 1 to 6 carbons or a phenylene group, and wherein Rf and Rf' are independently a fluorohydrocarbon group, a perfluorocarbon group, a fluorinated ether group or a fluorinated amine group.

- 11. The compound of Claim 10 wherein  $X^1$  is an allyl group, Br, Cl, F, I, or H, Rs is  $-CH_2CH_2-$ , and Rf is a perfluoroalkyl group.
- 12. The compound of Claim 10 wherein Rf is a fluorohydrocarbon group, a perfluorocarbon group, a fluorinated ether group or a fluorinated amine group.
- 13. The compound of Claim 10 wherein Rf is a linear perfluoroalkyl group of 3 to 20 carbons, a branched perfluoroalkyl group of 3 to 20 carbons, and a hydrofluoroalkyl group of 3 to 20 carbons, the

hydrofluoroalkyl group comprising up to one hydrogen atom for each two fluorine atoms.

14. The compound of Claim 10 wherein Rf is a linear perfluoroalkyl group of 6 to 12 carbons, a branched perfluoroalkyl group of 6 to 12 carbons, or a hydrofluoroalkyl group of 6 to 12 carbons, the hydrofluoroalkyl group comprising up to one hydrogen atom for each two fluorine atoms.

15. The compound of Claim 10 wherein R<sup>3</sup> is a perfluoroalkyl group.

- 16. The compound of Claim 10 wherein Rs is an alkylene group of 1 to 6 carbons.
  - 17. A chemical compound having the formula:

## $O=Sn[Rs(Rf)]_2$

wherein Rs is an alkylene group of 1 to 6 carbons or a phenylene group and wherein Rf is a fluorohydrocarbon group, a perfluorocarbon group, a fluorinated ether group or a fluorinated amine group.

18. The compound of Claim 17 wherein Rs is an alkylene group of 1 to 6 carbons.

put Bit

19. A chemical compound having the formula:

## $X^1X^2Sn[Rs(Rf)]_2$

wherein  $X^1$  and  $X^2$  are independently, the same or different, H,  $N_3$ ,  $OR^1$ ,  $OOR^1$   $SR^1$ ,  $SeR^1$ , CN, NC,  $NR^1R^2$ , a heteroaryl group, an alkyl group of 2 to 20 carbons, an alkenyl group, an alkynyl group,  $-C(0)R^3$ ,  $M((Rs')(Rf'))_3$ ,  $OM((Rs')(Rf'))_3$  or  $OOM((Rs')Rf'))_3$ , wherein M is Si, Ge, or Sn, and wherein  $R^1$  and  $R^2$  are each independently the same or different H, an alkyl group,  $-SO_2R^3$  or  $-C(O)R^3$ , wherein  $R^3$  is an alkyl group or an aryl group, wherein Rs and Rs' are each independently the same or different an alkylene group of 1 to 6 carbons or a phenylene group, and wherein Rf and Rf' are each independently a fluorohydrocarbon group, a perfluorocarbon group, a fluorinated ether group or a fluorinated amine group.

20. The compound of Claim 19 wherein Rs is an alkylene group of 1 to 6 carbons.